

on Vaccine Research

ABSTRACTS OF SUBMITTED PRESENTATIONS

- S25** CpG ODN is safe and highly effective in humans as adjuvant to HBV vaccine: Preliminary results of Phase I trial with CpG-ODN 7909
H. Davis^{1,2,3}, C.I. Cooper¹, M.L. Morris¹, S.M. Blitt¹, D.W. Cantmore¹, J. Heathcote¹

¹Coley Pharmaceutical Group, Wellesley, MA; ²Loeb Health Research Institute, Ottawa, Canada; ³University of Ottawa, Clinical Investigation Unit, Ottawa, Canada

¹Immunostimulatory CpG motifs are unmetabolized cytosine-phosphate dinucleotides common in bacterial DNA but rare in mammalian DNA. CpG DNA activates many immune cells. In animal models, CpG-containing oligodeoxynucleotides (CpG ODN) act as potent Th1-like adjuvants with many antigens. We identified a 24-mer containing 3 CpG motifs (CpG 7909) that strongly activates human cells *in vitro*. In this first report of using CpG 7909 adjuvants, we evaluated their safety, tolerability and immunogenicity of CpG 7909 as adjuvant to a hepatitis B virus (HBV) vaccine.

In a double-blind phase I study being carried out at two Canadian sites, healthy volunteers aged 18-35 years are vaccinated at 0, 1 and 5 mos by IM injection with Engix-B[®] (SmithKline Beecham), which contains 20µg yeast-derived hepatitis B surface antigen (HBsAg) adsorbed to alum, with or without adjuvant CpG 7909 (500µg). Interim group-wise comparisons shows immunogenicity of control vaccines, to be well tolerated, both locally and systemically. However, HBsAg-specific antibody responses (anti-HBs) are significantly better in CpG than control subjects. With control vaccine, no subjects ($n=8$, to date) had seroconverted (SC, anti-HBs > 1 mIU/ml) at 2 weeks post-prime and there was only 25% SC and 13% seroprotection (SP, > 10 mIU/ml) by 4 wk. In contrast, those receiving CpG 7909 with their vaccines ($n=12$) had 83 and 92% SC and 58 and 73% SP at 2 and 4 wk respectively. Post-boost (6 wk), SP was 57% for control subjects but 100% for subjects receiving CpG 7909. Anti-HBs titers were significantly higher with CpG, for all time points after prime and boost ($p<0.02$). We are also evaluating antibody isotype to identify the Th1/Th2 bias of responses.

These results indicate that CpG 7909 may allow the development of a 2- rather than a 3-dose HBV vaccine and that it may be a safe and highly effective adjuvant for a wide variety of human vaccines.

- S26** Using C3d to Increase the Immunogenicity of HIV-1 Envelope and Influenza Hemagglutinin in Mice.
Ted M. Ross¹, Yan Xu, R. A. Bright, and Harriet L. Robinson. Emory University. Yerkes Regional Primate Research Center. Atlanta, Georgia 30329 USA.

DNA immunogens were used to test the C3d component of complement as a molecular adjuvant for HIV-1 envelope (Env) or Influenza hemagglutinin (HA) immunogenicity. In normal immune responses, the attachment of C3d to an antigen (Ag) enhances both the initiation and maturation of Ag-specific antibody (Ab) responses. The HIV-1 Env has been a target for developing an effective vaccine, however, Env has proven to be a weak immunogen; raising antibodies that are slow to undergo affinity maturation. DNA constructs were produced which encoded soluble gp120 or soluble HA fused to C3d (gp120-3C3d or HA-3C3d). The working hypothesis was soluble fusion proteins would increase the immunogenicity of either Env or HA in DNA immunized mice. Our results in mice inoculated with plasmids expressing HA-3C3d had a 15 fold increase in total IgG to HA compared to mice inoculated with plasmids expressing HA only. The amount of anti-HA IgG raised in mice inoculated with HA-3C3d was similar to the amount of antibody raised in mice inoculated with the natural, transmembrane form of HA (tmHA). In mice inoculated with gp120-3C3d constructs, there was at least 32 fold increase in anti-Env IgG antibody levels compared to mice inoculated with gp120 only expressing constructs. The avidity of the anti-HA or anti-Env antibody was 20% higher in mice inoculated with C3d fusion plasmids than in HA or gp120 only inoculated mice. The level of HA titers were higher in HA-3C3d inoculated mice compared to mice inoculated with HA only. However, these HA titers were less than the levels seen in mice inoculated with tmHA expressing plasmids. In addition, there was a significant correlation between HA1 titer and protection in mice to challenge influenza virus. This work was funded in part by NIH/NIAD Award R21 AI4325-01 (to T.M.R.).

- S27** Evaluation of Carbohydrate-PADRE Peptide Conjugate Vaccine Technology
M.J. Newman¹, J. Alexander, M.-F. del Guercio, A. Maewal, R.W. Chomau, A. Sente
Eppimune Inc. 5820 Nancy Ridge Drive, San Diego CA 92121, USA

The PADRE technology (Pan HLA-DR Epitope) is based on a family of synthetic peptides that were designed to bind to the majority of common HLA-DR antigens with moderate to high affinity, and to present immunogenic, large or charged, amino acids to the T-cell receptors on CD4+ helper T-lymphocytes. In its peptide form, PADRE is highly immunogenic and capable of inducing proliferation of naive helper T-lymphocytes. Thus, it may be well suited for use as an adjuvant or carrier for carbohydrate vaccine antigens. To test this hypothesis, we constructed experimental glycoconjugate vaccines based on PADRE and bacterial carbohydrates and evaluated their immunogenicity in C57BL/6 mice. Included were the dodecasaccharides derived from *Salmonella typhimurium* lipopolysaccharide O-antigen and capsular polysaccharides from *Streptococcus pneumoniae*. All of the glycoconjugates induced higher titers antibody responses in mice than did carbohydrate-antigen alone. Different adjuvants, in combination with PADRE conjugates also proved useful and allowed for the modulation of the antibody isotype profile. Alum adjuvant supported predominantly IgG₁ responses whereas O-21 saponin adjuvant augmented IgG_{2b} IgG_{2c} tiers. The antibodies produced by these vaccines bound to target bacteria, suggesting that biologically relevant specificities were produced.

- S28** Novel Synthetic LPS Receptor Agonists as Vaccine Adjuvants

S. T. Ishizaka¹, M. Przetalak, J. Rose, P. McGuinness, W. Gavin, Z. Meng, B. deCosta, H. Zhang, H. Yang, J. Chow, S. Motokoa, A. Suganuma, A. Nault, D. Rossignol and L.D. Hawkins

Eltal Research Institute, 4 Corporate Dr., Andover, MA 01810

Objective: The immunologic analysis of simplified synthetic LPS receptor agonists screened for use as potential vaccine adjuvants.

Methods and Results: LPS receptor agonists consisting of a six-fatty acid chain system with a non-sugar head structure were synthesized. Human whole blood stimulated with the compounds produced IL-10 and TNF- α . Structural modifications result in a series of molecules with a range of cytokine stimulatory activity, all significantly attenuated compared to LPS. The compounds activated an NFkB-driven reporter gene in cells expressing TLR4/MD2 surface molecules.

When mixed in aqueous solution with protein or polysaccharide antigens, the agonists boosted serum antibody responses in mice ten-fold or more above unadjuvanted antigen. Antibody was largely of the IgG1 subclass, indicative of a Th2-associated cytokine pattern. Antibody responses against the adjuvant itself occurred at less than 5% frequency. The compounds also supported antibody responses to tetanus toxoid given intramuscularly. Nasal and vaginal IgA was elicited in the immunization regimen, in addition to circulating IgG. The compounds showed excellent local safety when given s.c. in mice. High doses resulted in circulating IL-10, but doses below 30 µg did not result in measurable serum IL-10 increases, although retaining their adjuvant effects.

Conclusions: Simplified LPS receptor agonists, which can be synthesized in high purity, have the potential to be efficacious, low-cost adjuvants.